

## **LAPAROSCOPIC SLEEVE GASTRECTOMY DISCHARGE INSTRUCTIONS**

When you are discharged from the hospital it is important to have help at home. We suggest either a family member or a friend stay with you for the first few days. **It is best to stay within one hour driving distance until after your first post-operative appointment.**

If you have any questions or concerns while at home you can always call Dr. Monash's office at (520) 319-6000. After 5:00 pm and on the weekend, calls will go directly to the answering service. Dr. Monash or Dr. Schwartz (whomever is the Bariatric Surgeon on call) will be contacted regarding your questions and/or concerns.

### **When to Call the Surgeon**

- Persistent nausea or vomiting lasting more than 4-6 hours which prevents the intake of any fluids.
- Post operative pain not relieved by the pain medication.
- Fever greater than 101 degrees F.
- Development of foul smelling drainage at staple or wound sites.
- If there are any changes in your recovery that concern you, give us a call.

### **Wound Care**

- Shower daily beginning two days after surgery.
- Let soap and water run over the incisions, scrub VERY gently, and pat dry.
- Do not submerge the incisions for two weeks (bathtub, swimming etc).
- Do not apply lotions, creams, or salves to the wound sites.
- The Steri-Strips will fall off on their own.
- Notify your surgeon if the wounds become red and swollen, develop a foul smelling drainage.

### **Activity**

- You must walk at least four times daily. Stop when you become fatigued. Increase your activity a little bit each day.
- Do not drive a car until you are completely off of the prescribed narcotic pain medication.
- You may walk up and down stairs. Take your time, and rest if you need to.
- Do not lift anything greater than 10 pounds until you have your first post-op visit. No strenuous activity for two weeks.
- Use your RIGHT arm for support when changing positions from lying to sitting to standing, and for any preferential weight bearing when needed.

### **Bowel Activity**

It is not imperative that you have a bowel movement before leaving the hospital. If you are having difficulty at home, over the counter Colace or similar stool softeners work well to combat the constipating effect of narcotic pain medications.

Importantly: A small amount of blood in your first one or two bowel movements is normal. If the bleeding persists beyond two bowel movements, call the office.

## Medication

***Medications taken by mouth must be crushed or taken in liquid form during the first week after surgery.***

### Standard Discharge Medications

- **Lortab Elixir** - 15 mL every 4-6 hours; take only as needed for operative pain relief. If you do not have pain, do not take Lortab. If you have mild pain or discomfort you may take Children's Tylenol (crushed) one tablet every 4-6 hours. If you were taking a narcotic for other pain prior to surgery, please call your primary care provider for pain medication adjustment.
- **Vitamin supplement** - 2 chewable multiple vitamins every day. Recommended chewable multivitamins are: Flintstones with minerals or Centrum Jr. with minerals.

### Pre-operative Medication

- You are instructed to resume your pre-operative medications. **Arrange an appointment with your primary care provider soon after hospital discharge to review these medications.**
- For as least one week post-operatively, **medications taken orally must be crushed or taken in liquid form.**

## Diet

Begin a protein-based liquid diet at home, and continue for two weeks. Please refer to the dietary instructions given in your Bariatric Booklet for details.

- Always drink slowly to avoid discomfort or vomiting. Start with 30 mL (one ounce) every 15 minutes. Increase the amount as tolerated.
- Stop drinking when you feel full. Allow 10-15 minutes for your stomach to empty.
- Your goal **NO LESS THAN** 60 grams of protein every day. Aim for 80 grams if possible.
- **KEEP HYDRATED.** Protein-based liquids and water intake **must** total at least 6 cups (48 ounces) every day.

## Follow-Up

Call the office (520) 319-6000 to schedule an appointment for 10 to 14 days following your discharge. The next appointments will be at three months, six months, and then yearly thereafter. You will be given the appropriate instructions to have blood taken before your follow-up appointments to help identify and supplement any vitamin and/or mineral deficiencies.